Name: _____ Age/DOB: _____ Date of Injury: _____

Post Concussion Symptom Scale

No symptoms"0"-------Moderate "3"---------Severe"6"

Time after Concussion

SYMPTOMS	Days/Hrs _____	Days/Hrs _____	Days/Hrs _____
Headache	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Nausea	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Vomiting	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Balance problems	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Dizziness	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Fatigue	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Trouble falling to sleep	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Excessive sleep	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Loss of sleep	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Drowsiness	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Light sensitivity	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Noise sensitivity	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Irritability	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Sadness	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Nervousness	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
More emotional	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Numbness	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Feeling "slow"	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Feeling "foggy"	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Difficulty concentrating	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Difficulty remembering	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
Visual problems	0 1 2 3 4 5 6	0 1 2 3 4 5 6	0 1 2 3 4 5 6
TOTAL SCORE	_____	_____	_____

Use of the Post-Concussion Symptom Scale: The athlete should fill out the form, on his or her own, in order to give a subjective value for each symptom. This form can be used with each encounter to track the athlete's progress towards the resolution of symptoms. Many athletes may have some of these reported symptoms at a baseline, such as concentration difficulties in the patient with attention-deficit disorder or sadness in an athlete with underlying depression, and must be taken into consideration when interpreting the score. Athletes do not have to be at a total score of zero to return to play if they already have had some symptoms prior to their concussion.